

Pet Doctors Veterinary Supplies and Services
Immunoassay Test Report

Pet Name: Ogie

Owner Name: Karen Ramos

Species: Canine

Gender: Male

Age: 10Year


Patient or Visit No.:

Sample ID: 202607280001

Time Tested: 28/07/2026 14:49:12

Test Result

Test	Result	Unit	Reference Range
SDMA	19.57 Abnormal	µg/dL	≤14 µg/dL Normal 14-19 µg/dL Suspected >19 µg/dL Abnormal



Clinical Significance

SDMA

Symmetric dimethylarginine (SDMA) is a methylated arginine amino acid. Most of SDMA is excreted by the kidneys. SDMA is an early biomarker of kidney function. It correlates well with glomerular filtration rate(GFR). SDMA tests diagnose chronic kidney disease (CKD) and acute kidney injury (AKI) much earlier than creatinine tests.

Veterinary Technician: not specified

Attending Veterinarian: Dr. Marian Wanagen

Date Reported: 30/07/2026

This report is solely for the received sample and serves as a diagnostic reference for veterinarians.



Hospital Tel.: +639171533903

Hospital Add.: McArthur Highway, San Miguel, Calasiao, Pangasinan



Pet Doctors Veterinary Supplies and Services

Chemistry Test Report

Pet Name: Ogie
Gender: Male
Sample ID: 2607280001

Owner Name: Karen Ramos
Age: 10 Years
Sample Type: Serum

Species: Canine
Patient or Visit No.:
Time Tested: 28/07/2026 14:23:22

Test Panel: Comprehensive 24 Profile

Test	Result	Unit	Reference Range	Low	Normal	High	
TP	74.6	g/L	52-82				Whole Body
ALB	32.6	g/L	22-44				Liver, Kidney
GLOB	42.0	g/L	20-52				Whole Body
ALB/GLOB	0.8						Calculated value
TBIL	4.09	umol/L	2-15				Whole Body, Liver
GGT	4.9	U/L	0-7				Liver
AST	21	U/L	8.9-55				Liver
ALT	127	U/L	10-140				Liver
AST/ALT	0.17						Calculated value
ALP	30	U/L	20-150				Liver
TBA	5.30	umol/L	0-20				Liver, Gallbladder
AMYL	911	U/L	200-1800				Pancreas
CK	74	U/L	20-200				Myocardium, Skeletal muscle
CREA	101	umol/L	27-149				Kidney
BUN	8.27	mmol/L	2.5-11.5				Kidney
BUN/CREA	20						Calculated value
GLU	5.82	mmol/L	3.89-7.95				Whole Body
CHOL	6.17	mmol/L	2.84-8.26				Whole Body
TRIG	0.84	mmol/L	0.1-0.9				
tCO2	20	mmol/L	12-27				Whole Body
Ca	2.46	mmol/L	1.98-2.95				Whole Body
PHOS	1.30	mmol/L	0.81-2.2				Kidney
Ca×P	40	mg/dL					
Mg	0.78	mmol/L	0.6-1.09				

Clinical Significance

TP	Reflects the protein content in the animal's body: dehydration, hepatic and renal diseases, or gastrointestinal diseases can cause this index to be abnormal.
ALB	Synthesized by the liver, the primary plasma protein, maintaining osmotic pressure and reflecting the body's nutritional status.
GLOB	Associated with chronic infections or autoimmune diseases.
ALB/GLOB	Low levels may indicate renal dysfunction, amyloidosis, inflammation, infections, enteritis, hepatic and renal diseases, or severe malnutrition due to malabsorption/digestion.
TBIL	Used to assess anemia and biliary tract diseases.
GGT	A hepatic enzyme indicator, mainly for assessing cholestasis.
AST	An indicator for liver and muscle diseases.
ALT	An indicator for diagnosing liver diseases in dogs and cats, significantly elevated in liver damage or liver diseases.
AST/ALT	Used to assess the degree of hepatocyte injury and the risk of primary liver cancer.
ALP	An indicator for diagnosing hepatobiliary diseases in animals, related to cholestasis; slight elevation in cats has clinical significance.
TBA	Hepatic function indicator
AMYL	An indicator of pancreatic disease, often measured with LPA (lipase) to assess hepatosplenomegaly.
CK	Indicator of muscle injury or neurological disease.
CREA	A metabolic product of muscle, excreted by the kidneys, may rise in renal disease or urethral obstruction.
BUN	Synthesized by the liver and excreted by the kidneys, one of the renal function indicators, may also rise due to other causes (diet, dehydration).
BUN/CREA	Seen in dehydration, gastrointestinal bleeding, renal disease, urinary tract obstruction.



Hospital Tel.: +639171533903

Hospital Add.: McArthur Highway, San Miguel, Calasiao, Pangasinan



GLU	Abnormal levels of blood glucose concentration may occur when the body is in poor condition or in the presence of diabetes.
CHOL	Elevated in liver/kidney diseases or endocrine abnormalities.
TRIG	Indicator of fat content in the blood.
tCO2	Assessment of acid-base balance.
Ca	Abnormal levels may be caused by various diseases such as improper lactation, nutritional disorders, and tumors.
PHOS	Useful for assessing parathyroid function, vitamin D deficiency, bone tumors, and multiple myeloma.
Ca×P	Used to assess bone metabolism. Note: Animals with sCaPP > 70 mg have been observed to have shorter survival periods, indicating poor prognosis of chronic renal disease. Note: Animals with CPP > 60-70 are at risk of soft tissue calcification, although increased concentrations may persist for several weeks to months.
Mg	An indicator of adrenal and renal function.

Veterinary Technician: not specified

Attending Veterinarian: Dr. Marian Wanagen

Date Reported: 30/07/2026

This report is solely for the received sample and serves as a diagnostic reference for veterinarians.



Hospital Tel.: +639171533903

Hospital Add.: McArthur Highway, San Miguel, Calasiao, Pangasinan

