

Client: Maramba, Ems (5071)  
Patient Name: Gigi  
Species: Canine  
Breed:

Gender: Male  
Weight: 19.35 kgs  
Age: 8 Months  
Doctor:

PET DOCTORS VETERINARY  
CLINIC  
Diversion Road, San Miguel,  
Calasiao, Pangasinan  
0932-278-9861 075-523-  
1751

Test	Results	Reference Interval	LOW	NORMAL	HIGH
------	---------	--------------------	-----	--------	------

Catalyst One (December 9, 2023 3:34 PM)

Progesterone 0.4 ng/mL

Progesterone (ng/mL)

< 1.0 Anestrus/Proestrus

1.0-1.9 Pre-LH surge

2.0-3.0 Associated with LH surge, about 24-48 hours prior to the ovulation wave

3.1-4.9 Post-LH surge, pre-ovulation

5.0-12.0 At or near ovulation (progesterone concentration may vary with breed and size)

Physiologic variation in progesterone occurs prior to ovulation. We recommend confirming a continuing rise in progesterone (every 24-48 hours until ovulation is confirmed) when evaluating for timing of breeding.

When measuring progesterone to evaluate for neuter status or for ovarian remnant in dogs (2 weeks-2 months post signs of estrus), a progesterone > 2 ng/mL is consistent with the presence of a functioning corpus luteum and the presence of ovarian tissue. In addition to progesterone concentrations, anti-mullerian hormone, LH concentrations, vaginal cytology, clinical signs, abdominal ultrasound and estrogen concentrations can be used to further evaluate neuter status and for ovarian remnants.