Client: Espino, Le Anne Denise (0067)

Patient Name: Georgie Species: Canine

Breed: Shih Tzu

Gender: Female Weight: 7.65 kgs Age: 13 Years

Doctor:

PET DOCTORS VETERINARY CLINIC Diversion Road, San Miguel, Calasiao, Pangasinan 0932-278-9861 075-523-

1751

Test Results Reference Interval LOW NORMAL HIGH

Catalyst One (March 9, 2024 4:31 PM)

TT4 2.3 μg/dL 1.0 - 4.0

Diagnostic Interpretation for TT4

< 1.0 µg/dL Low

 $1.0 - 2.0 \ \mu g/dL$  Low Normal  $1.0 - 4.0 \ \mu g/dL$  Normal  $> 4.0 \ \mu g/dL$  High  $2.1 - 5.4 \ \mu g/dL$  Therapeutic

Printed: March 9, 2024 4:32 PM

Dogs with no clinical signs of hypothyroidism and results within the normal reference range are likely euthyroid. Dogs with low T4 concentrations may be hypothyroid or "euthyroid sick". Occasionally, hypothyroid dogs can have T4 concentrations that are low normal. Dogs with clinical signs of hypothyroidism and low or low normal T4 concentrations may be evaluated further by submission of freeT4 (fT4) and canine TSH. A high T4 concentration in a clinically normal dog is likely a variation of normal; however elevations may occur secondary to thyroid auto antibodies or rarely thyroid neoplasia. For dogs on thyroid supplement, acceptable 4-6 hour post pill total T4 concentrations generally fall within the higher end or slightly above the reference range.

